Kanse Lipano (500 g)



Product Code: 6069080001					
Unit EAN	Shelf life (months)	Description	Units	Gross Weight (kg)	Measures in cm (W x L x H)
8 008698 032022	18	Unit	1 x 500g	0.62	10.3 x 10.3 x 14.9
		Box	6 Units/Box	3.889	22.6 x 33 x 17
		Pallet	12 Boxes/Layer 6 Layers 72 Boxes	301	120 x 80 x 116.5 (with pallet)

Definition and Indication: FOOD FOR SPECIAL MEDICAL PURPOSES.

For the dietary management of oxidation disorders of long chain fatty acids, fat maldigestion and fat malabsorption. Nutritionally complete, low fat, powdered formula, low in long-chain triglycerides (LCT) and high in medium-chain triglycerides (MCT). Lipano contains a dosed amount of vitamins, minerals, trace elements, docosahexaenoic acid (DHA) and arachidonic acid (AA).

Ingredients:

Maltodextrin, Vegetableoils 16.6% (MCT oil, **soybean** oil, antioxidants: to copherol rich natural extract, Ascorbyl palmitate), Calcium caseinate (**milk**), Lactose (**milk**), Fructo-oligosaccharides (FOS) 3.5%, di-Calcium phosphate, tri-Potassium citrate, Emulsifiers: soya lecithin, citric acid esters of mono- and diglycerides of fatty acids; galactooligosaccharides - GOS (**lactose**) 0.4%, Sodium hydrogen phosphate, Potassium chloride, Choline bitartrate, Magnesium carbonate, Vitamins (A, D, E, K, C, B1, B2, niacin, B6, folic acid, panthotenic acid, B12, biotine), Sodium chloride, Oil from Mortierella alpina, DHA- and EPA-rich oil from the microalgae Schizochytrium sp., L-Cystine, Inositol, Iron sulfate, Zinc sulfate, L-Carnitine, L-Tartrate, Copper sulfate, Manganese sulfate, Potassium iodide, Sodium selenite, Sodium molybdate.

Important notice:

Must only be used under medical supervision. Suitable as a sole source of nutrition in infants up to 1 year age and as a supplementary feeding for children from 1 year age and adults. Not for parenteral use. Not suitable for people not a affected by mentioned diseases.

Dosage:

The total daily amount depends on age, body weight and individual metabolic situation of the patient.

Contraindications:

Lipanoshouldnotbeusedifketoacidosis/acidosis,decompensatedlivercirrhosis, chronic kidney disease is suspected or in people with medium chain acyl-CoA dehydrogenase deficiency (MCADD).

Preparation:

Suitable to be taken orally or via an enteral feeding tube as determined by the clinician or dietitian. Standard dilution (15%): The suggested feed concentration is 90 ml of water plus 15 g Lipano.

Storage:

Unopened: Store in a cool and dry place.

Opened: Once opened, close the lid tightly, store it in a cool and dry place and use within 4 weeks.

Nutrition declaration	per 100 g	per portion (15 g)
Energy	1883 kJ 448 kcal	282 kJ 67 kcal
Fat of which saturates of which mono-unsaturates of which poly-unsaturates of which MCT of which LCT davon Linolsäure (Omega 6) of which α-linolenic acid (Omega 3) of which arachidonic acid (AHA) of which docosahexaenoic acid (DHA)	17.5 g 14.3 g 0.7 g 2.4 g 13.4 g 4.5 g 1.7 g 326 mg 70 mg 56 mg	2.6 g 2.2 g 0.1 g 0.4 g 2.0 g 0.7 g 0.3 g 49 mg 10 mg 8.4 mg
Carbohydrate of which sugar	62.9 g 8.1 g	9.4 g 1.2 g
Fibre	3.4 g	0.5 g
Protein	10.9 g	1.6 g
Salt of which sodium	0.32 g 129 mg 5.52 mmol	0.05 g 19 mg 0.83 mmol
Vitamins Vitamin A (RE) Vitamin D Vitamin E (α-tocopherol) Vitamin K Vitamin C Thiamin B1 Riboflavin B2 Niacin B3 Vitamin B6 Folat B9 (DFE) Vitamin B12 Biotin B8 Pantothenic acid B5	429 μg 12 μg 4.29 mg 69 mg 343 μg 686 μg 3.50 mg 429 μg 129 μg 0.86 μg 8.58 μg 3 mg	64 μg 1.78 μg 0.64 mg 2.63 μg 10 mg 51 μg 103 μg 0.53 mg 64 μg 19 μg 0.13 μg 1.29 μg 0.45 mg
Minerals Sodium Potassium Chloride Calcium Phosphorus Magnesium	129 mg 5.6 mmol 386 mg 9.9 mmol 283 mg 8.0 mmol 429 mg 10.7 mmol 344 mg 11.1 mmol 64 mg 2.6 mmol	19 mg 0.8 mmol 58 mg 1.5 mmol 42 mg 1.2 mmol 64 mg 1.6 mmol 52 mg 1.7 mmol 9.65 mg 0.4 mmol
Trace elements Iron Zinc Copper Manganese Selenium Molybdenum Iodine	4.88 mg 4.29 mg 300 μg 214 μg 15 μg 11 μg 73 μg	0.73 mg 0.64 mg 45 µg 32 µg 2.22 µg 1.65 µg 11 µg
Others L-Carnitine Choline Inositol	6.43 mg 132 mg 53 mg	0.96 mg 20 mg 7.95 mg